

American Associationof Equine Practitioners

Horse Health Education

LAMINITIS

Prevention and Treatment

The AAEP horse health presentations are endorsed by the AAEP in its original format.


The AAEP prohibits editing or personalizing them in any way.

www.aaep.org/horseowner



Horse Health Education: LAMINITIS

Every day, veterinarians across the country see hundreds of cases of laminitis, a painful disease that affects the feet of the horse. The alarming fact is that some cases are preventable. In many cases, it may be that we are killing our horses with kindness.





A common cause of laminitis is overfeeding – a management factor that is normally within our control.

By learning more about laminitis, its causes, signs and treatments, you may be able to minimize the risks of laminitis in your horse, or control longterm damage.



LAMINITIS DEFINED

Laminitis results from disruption (chronic, intermittent or short-term) of blood flow to the sensitive and insensitive laminae.




LAMINITIS DEFINED

The laminae structures within the foot secure the coffin bone (wedge-shaped bone within the foot) to the hoof wall.

Inflammation leads to scar tissue formation, which can permanently weaken the laminae and interfere with the wall/hoof wall bond.



LAMINITIS DEFINED



Front Limb: cross section-skeletal, ligaments, tendons



LAMINITIS DEFINED

The terms "laminitis" and "founder" are often used interchangeably.

Founder usually refers to a chronic (long-term) condition associated with rotation of the coffin bone.

Acute laminitis is usually defined as the first 72 hours at which after this point it is referred to as "chronic". Symptoms associated with an initial pathological insult, include pain and inflammation of the laminae.






LAMINITIS DEFINED

Laminitis can affect one or all of the feet, but it is most often seen in both front feet concurrently.



LAMINITIS DEFINED

In severe cases, the bone and the hoof wall can separate. In these situations, the coffin bone may rotate within the foot, be displaced downward, "sink" and eventually penetrate the sole. These changes are referred to as "founder."





CAUSES

While the exact mechanisms by which the feet are damaged remain mysteries, we know certain precipitating events can produce laminitis. Although laminitis manifests in the feet, the underlying cause is often a disturbance elsewhere in the horse's body.



CAUSES



Causes vary and may include the following:

- Digestive upsets due to grain overload (such as excessive grain, fruit or snacks) or abrupt changes in diet.
- Sudden access to excessive amounts of lush forage before the horse's system has had time to adapt; this type of laminitis is known as "grass founder."



CAUSES

- Toxins released within the horse's system.
- High fever or illness; any illness that causes high fever or serious metabolic disturbances has the potential to cause laminitis (ie. Potomac Horse Fever).
- Severe colic or diarrhea.
- Retained placenta in the mare after foaling.



CAUSES




- Excessive concussion to the feet, often referred to as "road founder."
- Support limb laminitis from excessive weight bearing on one leg due to injury of another leg, or any other alteration of the normal gait.
- Various primary foot diseases.
- Bedding that contains Black Walnut (Juglans nigra) shavings.
- It has been suggested but not proven that prolonged use or high doses of corticosteroids may contribute to the development of laminitis.




RISK FACTORS



Factors that seem to increase a horse's susceptibility to laminitis or increase the severity of the condition when it does occur include the following:






- High nutritional plane (feeding large amounts of carbohydrate-rich meals).
- Ponies, Morgans, donkeys and miniature horses.





RISK FACTORS




- Unrestricted grain binges, such as when a horse breaks into the feed room. (If this happens, DO NOT WAIT until symptoms develop to call your veterinarian. CALL IMMEDIATELY SO CORRECTIVE ACTION CAN BE TAKED BEFORE TISSUE DAMAGE PROGRESSES.)
- Horses that have had previous episodes of laminitis.
- Older horses with Cushing's disease.



Horse Health Education: LAMINITIS

SIGNS


Signs of acute laminitis include the following:



- Lameness, especially when a horse is turning in circles; shifting leg lameness when standing.
- Pain in the toe region when pressure is applied with hoof testers.
- Increased digital pulse felt in the foot/feet.




SIGNS



- Heat in the feet.
- Reluctant or hesitant gait, "walking on eggshells."
- A "sawhorse stance," with the front feet stretched out in front to alleviate pressure on the toes and the hind feet under them to support the weight their front feet cannot.



SIGNS



Signs of chronic laminitis may include the following:

- Rings in hoof wall that become wider as they are followed from toe to heel.
- Widened white line at the quarters and toes with occurrence of seromas (blood pockets) and/or abscesses (this is not to be confused with white line disease or seedy toe).




SIGNS



- Dropped soles or flat feet.
- Thick, "cresty" necked horses, a classic sign of metabolic disease, are often more prone to develop laminitis/founder, but not all cresty neck horses have had an episode of laminitis.



SIGNS



• Dished hooves, which are the result of unequal rates of hoof growth. (The heels grow at a faster rate that the rest of the hoof, resulting in an "Aladdin slipper" appearance.) This appearance can also arise from poor hoof care or white line disease, weakening the hoof capsule.



TREATMENT

The sooner treatment begins, the better the chance for recovery. Treatment will depend on specific circumstances, but may include the following:

- Diagnosing and treating the primary problem. (Laminitis is often due to a systemic or general problem elsewhere in the horse's body.)
- Icing the feet in the acute stages by placing the feet in a bucket of ice water to the level of the carpus.



TREATMENT



- Dietary restrictions: stop feeding all grain-based and pasture diets; feed only grass hay until advised otherwise by your veterinarian.
- Your veterinarian may treat the affected horse with mineral oil via nasogastric tube in cases of grain overload. The mineral oil may help to eliminate excessive grain from the digestive tract.
- Administration of non-steroidal antiinflammatories are essential for controlling pain and inflammation within the foot. Your veterinarian may also administer IV DMSO and other fluids if the horse is ill or dehydrated.



TREATMENT

• Your veterinarian may administer vasoactive medications such as antibiotics to fight infection and administer anti-endotoxins to reduce bacterial toxicity.

Corticosteroids are inadvisable for the treatment of laminitis as they have been implicated as a possible predisposing factor in the development of laminitis.



TREATMENT



- It may be recommended to stable the horse on soft ground, such as in sand or shavings (not black walnut), and to encourage the horse to lie down to reduce pressure on the weakened laminae.
- Sand is also used for the reason it provides superior frog and sole support that no man-made shoe can.



TREATMENT



- Open and drain any abscesses that may develop.
- Cooperation between your veterinarian and the farrier. (Techniques that may be helpful include corrective trimming, frog supports and therapeutic shoes or pads.)

It takes most horses 10 to 12 months to grow an entirely new hoof. Therefore, therapy may last up to a year or longer.



LONG-TERM OUTLOOK

Some horses that develop laminitis make uneventful recoveries and go on to lead long, useful lives.

Unfortunately, others suffer such severe, irreparable damage that they are, for humane reasons, euthanized.



LONG-TERM OUTLOOK



Your equine practitioner can provide you with information about your horse's condition based on serial (or repeated) radiographic studies (X-rays) and the animal's response to treatment.



LONG-TERM OUTLOOK



Radiographs will show how much rotation of the coffin bone has occurred. This will help you make a decision in the best interest of the horse and help the farrier with the therapeutic shoeing.



MANAGEMENT

Once a horse has had laminitis, it may be more likely to reoccur. In fact, a number of cases become chronic as a consequence of the coffin bone rotation within the foot and also for the reason that the laminae never regains its original strength.

There may also be interference with normal blood flow to the feet as well as metabolic changes within the horse.



MANAGEMENT



Extra care is recommended for any horse that has had laminitis, including:

• A modified diet that provides adequate nutrition based on high-quality forage; digestible fiber (beet pulp) and fat; and avoiding excess carbohydrates, especially from grain.



MANAGEMENT



- Routine hoof care, including regular trimming and, in some cases, therapeutic shoeing. (Additional radiographs may be necessary to monitor progress.)
- A good health maintenance schedule that includes parasite control and vaccinations to reduce the horse's susceptibility to illness or disease.



Horse Health Education: LAMINITIS

MANAGEMENT

- Determine if a nutritional supplement formulated to promote hoof health is necessary.
- Avoid grazing lush pastures, particularly between late morning and late afternoon hours as plant sugars are at their highest.
- Restrict pasture intake during the Spring season or anytime the pasture has sudden growth, or "greens up."



SUMMARY

The best way to avoid laminitis is to prevent causes that are under your control.

- Keep all grain stored securely out of the reach of horses.
- Introduce your horse to lush pasture gradually.
- Be aware that when a horse is ill, under stress, or overweight, it is especially at risk.



SUMMARY

Consult your equine veterinarian to formulate a good dietary plan, with a routine health and hoof care program.

Laminitis is a medical emergency. If you suspect your horse may be suffering from laminitis, call your veterinarian immediately.





Photos courtesy of

Dana Zimmel, DVM, DACVIM, DABVP University of Florida • Gainesville, FL



www.louriezipf.com louzieann@aol.com

specializing in equine portraits and sports images
 also available for editorial work

Loveland, CO 970.663.4322







The AAEP would like to thank our partners for their support of the AAEP's Owner Education Programs.

























EQUUS



MAKE AN IMPACT ON THE WELFARE OF HORSES EVERYWHERE

Join your veterinarian in support of the AAEP Foundation to Help Us Help Horses.

The AAEP Foundation's mission is to improve the health and welfare of the horse through the support of:

- Equine Research
- Education
- Benevolence
- The Equine Community

To make a donation or learn more about the AAEP Foundation, visit www.aaepfoundation.org





For more horse health information, visit our Web site:

www.aaep.org/horseowner